arXiv:0911.1706v1 [math.CA] 9 Nov 2009

L_p Convergence with Rates of Smooth Poisson-Cauchy Type Singular Operators

George A. Anastassiou & Razvan A. Mezei

Department of Mathematical Sciences
The University of Memphis
Memphis, TN 38152, U.S.A.
`ganastss@memphis.edu`
`rmezei@memphis.edu`

Abstract. In this article we continue the study of smooth Poisson-Cauchy Type singular integral operators on the line regarding their convergence to the unit operator with rates in the L_p norm, $p \geq 1$. The related established inequalities involve the higher order L_p modulus of smoothness of the engaged function or its higher order derivative.

AMS 2000 Mathematics Subject Classification: *Primary*: 41A17, 41A35; *Secondary*: 26D15

Key Words and Phrases: Poisson-Cauchy Type singular integral, modulus of smoothness, L_p convergence.

1 Introduction

The rate of convergence of singular integrals has been studied in [9], [13], [14], [15], [7], [8], [4], [5], [6] and these articles motivate our work. Here we study the L_p, $p \geq 1$, convergence of smooth Poisson-Cauchy Type singular integral operators over $\mathbb{R}$ to the unit operator with rates over smooth functions with higher order derivatives in $L_p(\mathbb{R})$. We establish related Jackson type inequalities involving the higher L_p modulus of smoothness of the engaged function or its higher order derivative. The discussed operators are not in general positive, see [10], [11]. Other motivation comes from [1], [2].

2 Results

In the next we introduce and deal with the *smooth Poisson-Cauchy Type singular integral operators* $M_{r,\xi}(f;x)$ defined as follows.

For $r \in \mathbb{N}$ and $n \in \mathbb{Z}_+$ we set

$$\alpha_j = \begin{cases} (-1)^{r-j}\binom{r}{j} j^{-n}, & j = 1, \ldots, r, \\ 1 - \sum\limits_{j=1}^{r} (-1)^{r-j}\binom{r}{j} j^{-n}, & j = 0, \end{cases} \tag{1}$$

that is $\sum\limits_{j=0}^{r} \alpha_j = 1$.

Let $f \in C^n(\mathbb{R})$ and $f^{(n)} \in L_p(\mathbb{R})$, $1 \leq p < \infty$, $\alpha \in \mathbb{N}$, $\beta > \frac{1}{2\alpha}$, we define for $x \in \mathbb{R}$, $\xi > 0$ the Lebesgue integral

$$M_{r,\xi}(f;x) = W \int_{-\infty}^{\infty} \frac{\sum_{j=0}^{r} \alpha_j f(x+jt)}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt, \tag{2}$$

where the constant is defined as

$$W = \frac{\Gamma\left(\beta\right)\alpha\xi^{2\alpha\beta-1}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}.$$

Note 1. The operators $M_{r,\xi}$ are not, in general, positive. See [10], (18).

We notice by $W\int_{-\infty}^{\infty} \frac{1}{(t^{2\alpha}+\xi^{2\alpha})^{\beta}} dt = 1$, that $M_{r,\xi}(c,x) = c$, c constant, see also [10], [11], and

$$M_{r,\xi}(f;x) - f(x) = W\left(\sum_{j=0}^{r} \alpha_j \int_{-\infty}^{\infty} \left[f(x+jt)-f(x)\right] \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt\right). \tag{3}$$

We use also that

$$\int_{-\infty}^{\infty} \frac{t^k}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt = \begin{cases} 0, & k \text{ odd}, \\ \frac{1}{\xi^{2\alpha\beta-k-1}\alpha} \frac{\Gamma\left(\frac{k+1}{2\alpha}\right)\Gamma\left(\beta-\frac{k+1}{2\alpha}\right)}{\Gamma(\beta)}, & k \text{ even, with } \beta > \frac{k+1}{2\alpha}, \end{cases} \tag{4}$$

see [16].

We need the rth L_p-modulus of smoothness

$$\omega_r(f^{(n)}, h)_p := \sup_{|t| \leq h} \|\Delta_t^r f^{(n)}(x)\|_{p,x}, \quad h > 0, \tag{5}$$

where

$$\Delta_t^r f^{(n)}(x) := \sum_{j=0}^{r} (-1)^{r-j} \binom{r}{j} f^{(n)}(x+jt), \tag{6}$$

see [12], p. 44. Here we have that $\omega_r(f^{(n)}, h)_p < \infty$, $h > 0$.

We need to introduce

$$\delta_k := \sum_{j=1}^{r} \alpha_j j^k, \quad k = 1, \ldots, n \in \mathbb{N}, \tag{7}$$

and denote by $\lfloor \cdot \rfloor$ the integral part. Call

$$\tau(w,x) := \sum_{j=0}^{r} \alpha_j j^n f^{(n)}(x+jw) - \delta_n f^{(n)}(x). \tag{8}$$

Notice also that

$$-\sum_{j=1}^{r} (-1)^{r-j} \binom{r}{j} = (-1)^r \binom{r}{0}. \tag{9}$$

According to [3], p. 306, [1], we get

$$\tau(w,x) = \Delta_w^r f^{(n)}(x). \tag{10}$$

Thus

$$\|\tau(w,x)\|_{p,x} \leq \omega_r(f^{(n)}, |w|)_p, \quad w \in \mathbb{R}. \tag{11}$$

Using Taylor's formula, and the appropriate change of variables, one has (see [6])

$$\sum_{j=0}^{r} \alpha_j [f(x+jt) - f(x)] = \sum_{k=1}^{n} \frac{f^{(k)}(x)}{k!} \delta_k t^k + \mathcal{R}_n(0,t,x), \tag{12}$$

where

$$\mathcal{R}_n(0,t,x) := \int_0^t \frac{(t-w)^{n-1}}{(n-1)!} \tau(w,x) dw, \quad n \in \mathbb{N}. \tag{13}$$

Using the above terminology we obtain for $\beta > \frac{2\lfloor \frac{n}{2} \rfloor + 1}{2\alpha}$ that

$$\Delta(x) := M_{r,\xi}(f;x) - f(x) - \sum_{m=1}^{\lfloor n/2 \rfloor} \frac{f^{(2m)}(x)\delta_{2m}}{(2m)!} \frac{\Gamma\left(\frac{2m+1}{2\alpha}\right)\Gamma\left(\beta - \frac{2m+1}{2\alpha}\right)}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)} \xi^{2m} = \mathcal{R}_n^*(x), \tag{14}$$

where

$$\mathcal{R}_n^*(x) := W \int_{-\infty}^{\infty} \mathcal{R}_n(0,t,x) \frac{1}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt, \quad n \in \mathbb{N}. \tag{15}$$

In $\Delta(x)$, see (14), the sum collapses when $n = 1$.

We present our first result.

Theorem 1. *Let* $p, q > 1$ *such that* $\frac{1}{p} + \frac{1}{q} = 1$, $n \in \mathbb{N}$, $\alpha \in \mathbb{N}$, $\beta > \frac{1}{\alpha}\left(\frac{1}{p} + n + r\right)$ *and the rest as above. Then*

$$\|\Delta(x)\|_p \leq \frac{(2\alpha)^{\frac{1}{p}} \Gamma(\beta) \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{\frac{1}{q}} \xi^n \tau^{\frac{1}{p}}}{\Gamma\left(\frac{q\beta}{2}\right)^{\frac{1}{q}} \Gamma\left(\frac{1}{2\alpha}\right)^{\frac{1}{p}} \Gamma\left(\beta - \frac{1}{2\alpha}\right) (rp+1)^{\frac{1}{p}} \left[(n-1)!\right] (q(n-1)+1)^{1/q}} \omega_r(f^{(n)}, \xi)_p, \tag{16}$$

where

$$0 < \tau := \left[\int_0^{\infty} (1+u)^{rp+1} \frac{u^{np-1}}{(u^{2\alpha}+1)^{p\beta/2}} du - \int_0^{\infty} \frac{u^{np-1}}{(u^{2\alpha}+1)^{p\beta/2}} du\right] < \infty. \tag{17}$$

Hence as $\xi \to 0$ *we obtain* $\|\Delta(x)\|_p \to 0$.

If additionally $f^{(2m)} \in L_p(\mathbb{R}), m = 1, 2, \ldots, \left\lfloor \frac{n}{2} \right\rfloor$ then $\left\|M_{r,\xi}(f) - f\right\|_p \to 0$, as $\xi \to 0$.

Proof. We observe that

$$\begin{aligned} |\Delta(x)|^p &= W^p \left| \int_{-\infty}^{\infty} \mathcal{R}_n(0,t,x) \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt \right|^p \\ &\leq W^p \left(\int_{-\infty}^{\infty} \left|\mathcal{R}_n(0,t,x)\right| \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt \right)^p \\ &\leq W^p \left(\int_{-\infty}^{\infty} \left| \int_0^{|t|} \frac{(|t|-w)^{n-1}}{(n-1)!} |\tau(sign(t) \cdot w, x)| dw \right| \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt \right)^p. \end{aligned} \tag{18}$$

Hence we have

$$I := \int_{-\infty}^{\infty} |\Delta(x)|^p dx \leq W^p \left(\int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \gamma(t,x) \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}} dt \right)^p dx \right), \tag{19}$$

where

$$\gamma(t,x) := \int_0^{|t|} \frac{(|t|-w)^{n-1}}{(n-1)!} |\tau(sign(t) \cdot w, x)| dw \geq 0. \tag{20}$$

Therefore by using Hölder's inequality suitably we obtain

$$\begin{aligned}
R.H.S.(19) &= W^p\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\gamma(t,x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)^p dx\right)\\
&= W^p\cdot\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\gamma(t,x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}dt\right)^p dx\right)\\
&\le W^p\cdot\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left[\gamma(t,x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right]^p dt\right)\left(\int_{-\infty}^{\infty}\left[\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right]^q dt\right)^{\frac{p}{q}} dx\right)
\end{aligned}$$

$$\begin{aligned}
&= W^p\cdot\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\gamma^p(t,x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)\left(\int_{-\infty}^{\infty}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{q\beta/2}}dt\right)^{\frac{p}{q}} dx\right)\\
&= W^p\cdot\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\gamma^p(t,x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)dx\right)\left(\frac{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)}{\Gamma\left(\frac{q\beta}{2}\right)\alpha\xi^{q\alpha\beta-1}}\right)^{\frac{p}{q}}\\
&= \frac{\xi^{p\alpha\beta-1}\alpha\left[\Gamma\left(\beta\right)\right]^p\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p}\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\gamma^p(t,x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)dx\right). \qquad (21)
\end{aligned}$$

Again by Hölder's inequality we have

$$\gamma^p(t,x)\le\frac{\left(\int_0^{|t|}|\tau(sign(t)\cdot w,x)|^p dw\right)}{((n-1)!)^p}\frac{|t|^{np-1}}{(q(n-1)+1)^{p/q}}. \qquad (22)$$

Consequently we have

$$\begin{aligned}
R.H.S.(21) &\le \frac{\xi^{p\alpha\beta-1}\alpha\left[\Gamma\left(\beta\right)\right]^p\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p}\\
&\quad\cdot\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\frac{\left(\int_0^{|t|}|\tau(sign(t)\cdot w,x)|^p dw\right)}{((n-1)!)^p}\frac{|t|^{np-1}}{(q(n-1)+1)^{p/q}}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)dx\right)\\
&=:(*),
\end{aligned}$$

(calling

$$c_1:=\frac{\xi^{p\alpha\beta-1}\alpha\left[\Gamma\left(\beta\right)\right]^p\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p((n-1)!)^p(q(n-1)+1)^{p/q}}) \qquad (23)$$

and

$$\begin{aligned}(*) &= c_1\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}|\tau(sign(t)\cdot w,x)|^p dw\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dx\right)dt\right)\\
&= c_1\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}|\Delta^r_{sign(t)\cdot w}f^{(n)}(x))|^p dw\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dx\right)dt\right)\\
&= c_1\left(\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}|\Delta^r_{sign(t)\cdot w}f^{(n)}(x))|^p dw\right)dx\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)\\
&= c_1\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}\left(\int_{-\infty}^{\infty}|\Delta^r_{sign(t)\cdot w}f^{(n)}(x))|^p dx\right)dw\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)\\
&\le c_1\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}\omega_r(f^{(n)},w)_p^p dw\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right). \qquad (24)\end{aligned}$$

So far we have proved

$$I \le c_1\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}\omega_r(f^{(n)},w)_p^p dw\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right). \qquad (25)$$

By [12], p. 45 we have

$$(R.H.S.(25)) \le c_1\left(\omega_r(f^{(n)},\xi)_p\right)^p\left(\int_{-\infty}^{\infty}\left(\int_0^{|t|}\left(1+\frac{w}{\xi}\right)^{rp}dw\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right) =: (**). \qquad (26)$$

But we see that

$$(**) = \left(\frac{\xi c_1}{rp+1}\right)\left(\omega_r(f^{(n)},\xi)_p\right)^p \mathcal{J}, \qquad (27)$$

where

$$\begin{aligned}\mathcal{J} &= \int_{-\infty}^{\infty}\left(\left(1+\frac{|t|}{\xi}\right)^{rp+1}-1\right)|t|^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\\
&= 2\int_0^{\infty}\left(\left(1+\frac{t}{\xi}\right)^{rp+1}-1\right)t^{np-1}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt. \qquad (28)\end{aligned}$$

Here we find

$$\begin{aligned}\mathcal{J} &= 2\xi^{p(n-\alpha\beta)}\int_0^{\infty}\left((1+u)^{rp+1}-1\right)u^{np-1}\frac{1}{(u^{2\alpha}+1)^{p\beta/2}}du\\
&= 2\xi^{p(n-\alpha\beta)}\left[\int_0^{\infty}(1+u)^{rp+1}\ \frac{u^{np-1}}{(u^{2\alpha}+1)^{p\beta/2}}du - \int_0^{\infty}\ \frac{u^{np-1}}{(u^{2\alpha}+1)^{p\beta/2}}du\right]. \qquad (29)\end{aligned}$$

Thus by (17) and (29) we obtain

$$\mathcal{J} = 2\xi^{p(n-\alpha\beta)}\tau. \qquad (30)$$

We notice that

$$\begin{aligned}
0 < \tau &< \int_0^\infty \frac{(1+u)^{rp+1}\, u^{np-1}}{(u^{2\alpha}+1)^{p\beta/2}} du \\
&< \int_0^\infty \frac{(1+u)^{rp+1}\, (1+u)^{np-1}}{(u^{2\alpha}+1)^{p\beta/2}} du \\
&= \int_0^\infty \frac{(1+u)^{p(n+r)}}{(u^{2\alpha}+1)^{p\beta/2}} du =: I_1.
\end{aligned}$$

Also call

$$K := \int_0^1 \frac{(1+u)^{p(n+r)}}{(u^{2\alpha}+1)^{p\beta/2}} du < \infty.$$

Then we can write

$$I_1 = K + \int_1^\infty \frac{(1+u)^{p(n+r)}}{(u^{2\alpha}+1)^{p\beta/2}} du < K + 2^{p(n+r)} \int_1^\infty \frac{u^{p(n+r)}}{(u^{2\alpha}+1)^{p\beta/2}} du = K + 2^{p(n+r)} I_2,$$

where $I_2 := \int_1^\infty \frac{u^{p(n+r)}}{(u^{2\alpha}+1)^{p\beta/2}} du$.

Since $\frac{1}{1+u^{2\alpha}} < \frac{1}{u^{2\alpha}}$, we have $\frac{1}{(1+u^{2\alpha})^{p\beta/2}} < \frac{1}{u^{p\alpha\beta}}$, for $u \in [1, \infty)$.

So we get

$$\begin{aligned}
I_2 &< \int_1^\infty u^{p(n+r-\alpha\beta)} du = \lim_{\varepsilon\to\infty} \int_1^\varepsilon u^{p(n+r-\alpha\beta)} du \\
&= \lim_{\varepsilon\to\infty} \left(\frac{\varepsilon^{p(n+r-\alpha\beta)+1} - 1}{p(n+r-\alpha\beta)+1} \right) = \frac{-1}{p(n+r-\alpha\beta)+1},
\end{aligned}$$

which is a positive number since $\beta > \frac{1}{\alpha}\left(\frac{1}{p} + n + r\right)$.

Consequently I_2 is finite, so is I_1, proving $\tau < \infty$.

Using (27) and (30) we get

$$\begin{aligned}
(**) &= \left(\frac{\xi c_1}{rp+1}\right) \left(\omega_r(f^{(n)}, \xi)_p\right)^p 2\xi^{p(n-\alpha\beta)} \tau \\
&= \frac{2\alpha \left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{\frac{p}{q}} \tau}{(rp+1)\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}} ((n-1)!)^p (q(n-1)+1)^{p/q}} \xi^{pn} \left(\omega_r(f^{(n)}, \xi)_p\right)^p. \quad (31)
\end{aligned}$$

I.e. we have established that

$$I \le \frac{2\alpha \left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{\frac{p}{q}} \tau}{(rp+1)\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}} ((n-1)!)^p (q(n-1)+1)^{p/q}} \xi^{pn} \left(\omega_r(f^{(n)}, \xi)_p\right)^p. \quad (32)$$

That is finishing the proof of the theorem. ■

The counterpart of Theorem 1 follows, case of $p = 1$.

Theorem 2. *Let* $f \in C^n(\mathbb{R})$ and $f^{(n)} \in L_1(\mathbb{R}), n \in \mathbb{N}$, $\alpha \in \mathbb{N}$, $\beta > \frac{n+r+1}{2\alpha}$. *Then*

$$\begin{aligned}
\|\Delta(x)\|_1 &\le \frac{1}{(r+1)(n-1)!\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)} \quad (33) \\
&\cdot \left[\sum_{k=1}^{r+1} \binom{r+1}{k} \Gamma\left(\frac{n+k}{2\alpha}\right) \Gamma\left(\beta - \frac{n+k}{2\alpha}\right)\right] \omega_r(f^{(n)}, \xi)_1 \xi^n.
\end{aligned}$$

Hence as $\xi \to 0$ *we obtain* $\|\Delta(x)\|_1 \to 0$.

If additionally $f^{(2m)} \in L_1(\mathbb{R}), m = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$ then $\|M_{r,\xi}(f) - f\|_1 \to 0$, as $\xi \to 0$.

Proof. It follows

$$\begin{aligned} |\Delta(x)| &= W \left| \int_{-\infty}^{\infty} \mathcal{R}_n(0,t,x) \frac{1}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right| \\ &\leq W \int_{-\infty}^{\infty} |\mathcal{R}_n(0,t,x)| \frac{1}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \\ &\leq W \int_{-\infty}^{\infty} \left(\int_0^{|t|} \frac{(|t|-w)^{n-1}}{(n-1)!} |\tau(sign(t) \cdot w, x)| dw \right) \frac{1}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt. \end{aligned} \tag{34}$$

Thus

$$\|\Delta(x)\|_1 = \int_{-\infty}^{\infty} |\Delta(x)| dx \leq W \cdot \tag{35}$$

$$\int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \left(\int_0^{|t|} \frac{(|t|-w)^{n-1}}{(n-1)!} |\tau(sign(t) \cdot w, x)| dw \right) \frac{1}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right) dx$$

$$=: (*)$$

But we see that

$$\int_0^{|t|} \frac{(|t|-w)^{n-1}}{(n-1)!} |\tau(sign(t) \cdot w, x)| dw \leq \frac{|t|^{n-1}}{(n-1)!} \int_0^{|t|} |\tau(sign(t) \cdot w, x)| dw. \tag{36}$$

Therefore it holds

$$\begin{aligned} (*) &\leq W \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \left(\frac{|t|^{n-1}}{(n-1)!} \int_0^{|t|} |\tau(sign(t) \cdot w, x)| dw \right) \frac{1}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right) dx \\ &= \frac{W}{(n-1)!} \left(\int_{-\infty}^{\infty} \left(\int_0^{|t|} \left(\int_{-\infty}^{\infty} |\tau(sign(t) \cdot w, x)| dx \right) dw \right) \frac{|t|^{n-1}}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right) \\ &\leq \frac{W}{(n-1)!} \left(\int_{-\infty}^{\infty} \left(\int_0^{|t|} \omega_r(f^{(n)}, w)_1 dw \right) \frac{|t|^{n-1}}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right). \end{aligned} \tag{37}$$

I.e. we get

$$\|\Delta(x)\|_1 \leq \frac{W}{(n-1)!} \left(\int_{-\infty}^{\infty} \left(\int_0^{|t|} \omega_r(f^{(n)}, w)_1 dw \right) \frac{|t|^{n-1}}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right). \tag{38}$$

Consequently we have

$$\begin{aligned} \|\Delta(x)\|_1 &\leq \frac{W \omega_r(f^{(n)}, \xi)_1}{(n-1)!} \left(\int_{-\infty}^{\infty} \left(\int_0^{|t|} \left(1 + \frac{w}{\xi}\right)^r dw \right) \frac{|t|^{n-1}}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right) \\ &= \frac{2\xi W \omega_r(f^{(n)}, \xi)_1}{(r+1)(n-1)!} \left(\int_0^{\infty} \left(\left(1 + \frac{t}{\xi}\right)^{r+1} - 1 \right) \frac{t^{n-1}}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right) \\ &= \frac{2\Gamma(\beta)\, \alpha \xi^{2\alpha\beta} \omega_r(f^{(n)}, \xi)_1}{(r+1)(n-1)!\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)} \left(\int_0^{\infty} \left(\left(1 + \frac{t}{\xi}\right)^{r+1} - 1 \right) \frac{t^{n-1}}{\left(t^{2\alpha} + \xi^{2\alpha}\right)^{\beta}} dt \right). \end{aligned} \tag{39}$$

We have gotten so far

$$\|\Delta(x)\|_1 \le \frac{2\Gamma(\beta)\,\alpha\xi^{2\alpha\beta}\omega_r(f^{(n)},\xi)_1 \cdot \lambda}{(r+1)\,(n-1)!\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}, \tag{40}$$

where

$$\lambda := \int_0^\infty \left(\left(1+\frac{t}{\xi}\right)^{r+1} - 1\right)\frac{t^{n-1}}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt. \tag{41}$$

One easily finds that

$$\begin{aligned}\lambda &= \int_0^\infty \left(\sum_{k=1}^{r+1}\binom{r+1}{k}\left(\frac{t}{\xi}\right)^k\right)\frac{t^{n-1}}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt \\ &= \xi^{n-2\alpha\beta}\sum_{k=1}^{r+1}\binom{r+1}{k}\int_0^\infty \frac{T^{n+k-1}}{\left(T^{2\alpha}+1\right)^{\beta}}dT \\ &= \xi^{n-2\alpha\beta}\sum_{k=1}^{r+1}\binom{r+1}{k}K_{n+k}. \end{aligned} \tag{42}$$

Where

$$K_{n+k} := \int_0^\infty \frac{T^{n+k-1}}{\left(T^{2\alpha}+1\right)^{\beta}}dT = \frac{\Gamma\left(\frac{n+k}{2\alpha}\right)\Gamma\left(\beta-\frac{n+k}{2\alpha}\right)}{\Gamma(\beta)\,2\alpha}. \tag{43}$$

$$\|\Delta(x)\|_1 \le \frac{1}{(r+1)\,(n-1)!\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}\left[\sum_{k=1}^{r+1}\binom{r+1}{k}\Gamma\left(\frac{n+k}{2\alpha}\right)\Gamma\left(\beta-\frac{n+k}{2\alpha}\right)\right]\omega_r(f^{(n)},\xi)_1\xi^n.$$

We have proved (33). ■

The case $n = 0$ is met next.

Proposition 1. *Let* $p, q > 1$ *such that* $\frac{1}{p}+\frac{1}{q} = 1$, $\alpha \in \mathbb{N}$, $\beta > \frac{1}{\alpha}\left(r+\frac{1}{p}\right)$ *and the rest as above. Then*

$$\|M_{r,\xi}(f) - f\|_p \le \frac{(2\alpha)^{\frac{1}{p}}\left[\Gamma(\beta)\right]\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{1}{q}}\theta^{\frac{1}{p}}}{\Gamma\left(\frac{1}{2\alpha}\right)^{\frac{1}{p}}\Gamma\left(\beta-\frac{1}{2\alpha}\right)\Gamma\left(\frac{q\beta}{2}\right)^{\frac{1}{q}}}\omega_r(f,\xi)_p, \tag{44}$$

where

$$0 < \theta := \int_0^\infty (1+t)^{rp}\frac{1}{\left(t^{2\alpha}+1\right)^{p\beta/2}}dt < \infty. \tag{45}$$

Hence as $\xi \to 0$ we obtain $M_{r,\xi} \to$ unit operator I in the L_p norm, $p > 1$.

Proof. By (3) we notice that,

$$\begin{aligned} M_{r,\xi}(f;x) - f(x) &= W\left(\sum_{j=0}^{r}\alpha_j\int_{-\infty}^{\infty}(f(x+jt)-f(x))\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right) \\ &= W\left(\int_{-\infty}^{\infty}\left(\sum_{j=0}^{r}\alpha_j\left(f(x+jt)-f(x)\right)\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right) \\ &= W\left(\int_{-\infty}^{\infty}\left(\sum_{j=1}^{r}\alpha_j f(x+jt) - \sum_{j=1}^{r}\alpha_j f(x)\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right) \end{aligned}$$

$$
\begin{aligned}
&= W\left(\int_{-\infty}^{\infty}\left(\sum_{j=1}^{r}\left((-1)^{r-j}\binom{r}{j}j^{-n}\right)f(x+jt)-\sum_{j=1}^{r}\left((-1)^{r-j}\binom{r}{j}j^{-n}\right)f(x)\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)\\
&= W\left(\int_{-\infty}^{\infty}\left(\sum_{j=1}^{r}\left((-1)^{r-j}\binom{r}{j}\right)f(x+jt)-\sum_{j=1}^{r}\left((-1)^{r-j}\binom{r}{j}\right)f(x)\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)\\
&\overset{(9)}{=} W\left(\int_{-\infty}^{\infty}\left(\sum_{j=1}^{r}\left((-1)^{r-j}\binom{r}{j}\right)f(x+jt)+\left((-1)^{r-0}\binom{r}{0}\right)f(x+0t)\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)\\
&= W\left(\int_{-\infty}^{\infty}\left(\sum_{j=0}^{r}(-1)^{r-j}\binom{r}{j}f(x+jt)\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)\\
&\overset{(6)}{=} W\left(\int_{-\infty}^{\infty}\left(\Delta_t^r f\right)(x)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right). \qquad (46)
\end{aligned}
$$

And then

$$
\left|M_{r,\xi}(f;x)-f(x)\right| \le W\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right). \qquad (47)
$$

We next estimate

$$
\begin{aligned}
&\int_{-\infty}^{\infty}\left|M_{r,\xi}(f;x)-f(x)\right|^{p}dx \le \int_{-\infty}^{\infty}(W)^{p}\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)^{p}dx\\
&= (W)^{p}\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left(\left|\Delta_t^r f(x)\right|\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right)\left(\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right)dt\right)^{p}dx\\
&\le (W)^{p}\int_{-\infty}^{\infty}\left(\left(\int_{-\infty}^{\infty}\left(\left|\Delta_t^r f(x)\right|\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right)^{p}dt\right)^{\frac{1}{p}}\left(\int_{-\infty}^{\infty}\left(\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right)^{q}dt\right)^{\frac{1}{q}}\right)^{p}dx
\end{aligned}
$$

$$
\begin{aligned}
&= (W)^{p}\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|^{p}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)\left(\int_{-\infty}^{\infty}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{q\beta/2}}dt\right)^{\frac{p}{q}}dx\\
&= (W)^{p}\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|^{p}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)\left(\frac{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)}{\Gamma\left(\frac{q\beta}{2}\right)\alpha\xi^{q\alpha\beta-1}}\right)^{\frac{p}{q}}dx\\
&= (W)^{p}\left(\frac{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)}{\Gamma\left(\frac{q\beta}{2}\right)\alpha\xi^{q\alpha\beta-1}}\right)^{\frac{p}{q}}\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|^{p}\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt\right)dx
\end{aligned}
$$

$$
\begin{aligned}
&= \frac{\alpha\xi^{\alpha\beta p-1}\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}} \int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|^p \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dx\right)dt \\
&= \frac{\alpha\xi^{\alpha\beta p-1}\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}} \int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}\left|\Delta_t^r f(x)\right|^p dx\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt \\
&\leq \frac{\alpha\xi^{\alpha\beta p-1}\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}} \int_{-\infty}^{\infty}\omega_r(f,|t|)_p^p \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt
\end{aligned}
$$

$$
\begin{aligned}
&= \frac{2\alpha\xi^{\alpha\beta p-1}\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}} \int_{0}^{\infty}\omega_r(f,t)_p^p \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt \\
&\leq \frac{2\alpha\xi^{\alpha\beta p-1}\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}}\omega_r(f,\xi)_p^p \int_{0}^{\infty}\left(1+\frac{t}{\xi}\right)^{rp} \frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}}dt \\
&= \frac{2\alpha\xi^{\alpha\beta p-1}\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}}\omega_r(f,\xi)_p^p \int_{0}^{\infty}(1+T)^{rp} \frac{1}{\left(T^{2\alpha}+1\right)^{p\beta/2}\xi^{\alpha p\beta}}\xi dT \\
&= \frac{2\alpha\left[\Gamma(\beta)\right]^p \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{\frac{p}{q}}}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{\frac{p}{q}}}\omega_r(f,\xi)_p^p \int_{0}^{\infty}(1+t)^{rp} \frac{1}{\left(t^{2\alpha}+1\right)^{p\beta/2}}dt. \qquad (48)
\end{aligned}
$$

We have established (44).

We also notice that

$$
\begin{aligned}
\theta &= \int_0^\infty \frac{(1+t)^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt = \int_0^1 \frac{(1+t)^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt + \int_1^\infty \frac{(1+t)^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt \\
&< \int_0^1 \frac{(1+t)^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt + 2^{rp}\int_1^\infty \frac{t^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt \\
&< \int_0^1 \frac{(1+t)^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt + 2^{rp}\int_1^\infty t^{p(r-\alpha\beta)}dt \\
&= \int_0^1 \frac{(1+t)^{rp}}{\left(1+t^{2\alpha}\right)^{p\beta/2}}dt - \frac{2^{rp}}{p(r-\alpha\beta)+1},
\end{aligned}
$$

the last, since $\beta > \frac{1}{\alpha}\left(r+\frac{1}{p}\right)$, is a finite positive constant. Thus $0<\theta<\infty$. ■

We also give

Proposition 2. *Assume* $\beta > \frac{r+1}{2\alpha}$. It holds

$$
\|M_{r,\xi}f - f\|_1 \leq \frac{2\alpha\Gamma(\beta)}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}\left(\int_0^\infty (1+t)^r \frac{1}{\left(t^{2\alpha}+1\right)^{\beta}}dt\right)\omega_r(f,\xi)_1. \qquad (49)
$$

Hence as $\xi \to 0$ *we get* $M_{r,\xi} \to I$ *in the* L_1 *norm.*

Proof. By (47) we have again

$$|M_{r,\xi}(f;x)-f(x)| \leq W\left(\int_{-\infty}^{\infty}|\Delta_t^r f(x)|\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right).$$

Next we estimate

$$\begin{aligned}
\int_{-\infty}^{\infty}|M_{r,\xi}(f;x)-f(x)|\,dx \leq{}& W\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}|\Delta_t^r f(x)|\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\right)dx\\
={}& W\int_{-\infty}^{\infty}\left(\int_{-\infty}^{\infty}|\Delta_t^r f(x)|\,dx\right)\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\\
\leq{}& W\int_{-\infty}^{\infty}\omega_r(f,|t|)_1\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\\
\leq{}& W2\omega_r(f,\xi)_1\int_0^{\infty}\left(1+\frac{t}{\xi}\right)^r\frac{1}{\left(t^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dt\\
={}& \frac{\Gamma(\beta)\,\xi^{2\alpha\beta-1}2\alpha}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}\omega_r(f,\xi)_1\int_0^{\infty}\xi\,(1+t)^r\frac{1}{\left(t^{2\alpha}+1\right)^{\beta}\xi^{2\alpha\beta}}dt\\
={}& \frac{\Gamma(\beta)\,2\alpha}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}\omega_r(f,\xi)_1\int_0^{\infty}(1+t)^r\frac{1}{\left(t^{2\alpha}+1\right)^{\beta}}dt. \qquad (50)
\end{aligned}$$

We have proved (49).

We also notice that

$$\begin{aligned}
0 &< \int_0^{\infty}(1+t)^r\frac{1}{\left(t^{2\alpha}+1\right)^{\beta}}dt\\
&= \int_0^1\frac{(1+t)^r}{\left(t^{2\alpha}+1\right)^{\beta}}dt+\int_1^{\infty}\frac{(1+t)^r}{\left(t^{2\alpha}+1\right)^{\beta}}dt\\
&< \int_0^1\frac{(1+t)^r}{\left(t^{2\alpha}+1\right)^{\beta}}dt+2^r\int_1^{\infty}t^{r-2\alpha\beta}dt\\
&= \int_0^1\frac{(1+t)^r}{\left(t^{2\alpha}+1\right)^{\beta}}dt-\frac{2^r}{(r-2\alpha\beta+1)},
\end{aligned}$$

which is a positive finite constant. ■

In the next we consider $f\in C^n(\mathbb{R})$ and $f^{(n)}\in L_p(\mathbb{R}), n=0$ or $n\geq 2$ even, $1\leq p<\infty$ and the similar *smooth singular operator of symmetric convolution type*

$$M_{\xi}(f;x)=W\int_{-\infty}^{\infty}f(x+y)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dy,\quad \text{for all } x\in\mathbb{R},\ \xi>0. \qquad (51)$$

That is

$$M_{\xi}(f;x)=W\int_0^{\infty}(f(x+y)+f(x-y))\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dy,$$

for all $x\in\mathbb{R}$, $\xi>0$. Notice that $M_{1,\xi}=M_{\xi}$. Let the central second order difference

$$(\tilde{\Delta}_y^2 f)(x):=f(x+y)+f(x-y)-2f(x). \qquad (52)$$

Notice that

$$(\tilde{\Delta}_{-y}^2 f)(x)=(\tilde{\Delta}_y^2 f)(x).$$

When $n \geq 2$ even using Taylor's formula with Cauchy remainder we eventually find

$$(\tilde{\Delta}_y^2 f)(x) = 2\sum_{\rho=1}^{n/2} \frac{f^{(2\rho)}(x)}{(2\rho)!} y^{2\rho} + \mathcal{R}_1(x), \tag{53}$$

where

$$\mathcal{R}_1(x) := \int_0^y (\tilde{\Delta}_t^2 f^{(n)})(x) \frac{(y-t)^{n-1}}{(n-1)!} dt. \tag{54}$$

Notice that

$$M_\xi(f;x) - f(x) = W \int_0^\infty (\tilde{\Delta}_y^2 f(x)) \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^\beta} dy. \tag{55}$$

Furthermore by (4), (53) and (55) we easily see that

$$\begin{aligned} K(x) \ : &= M_\xi(f;x) - f(x) - \sum_{\rho=1}^{n/2} \frac{f^{(2\rho)}(x)}{(2\rho)!} \frac{\Gamma\left(\frac{2\rho+1}{2\alpha}\right)\Gamma\left(\beta - \frac{2\rho+1}{2\alpha}\right)}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)} \xi^{2\rho} \\ &= W \int_0^\infty \left[\int_0^y (\tilde{\Delta}_t^2 f^{(n)})(x) \frac{(y-t)^{n-1}}{(n-1)!} dt\right] \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^\beta} dy, \end{aligned} \tag{56}$$

where $\beta > \frac{(n+1)}{2\alpha}$.

Therefore we have

$$|K(x)| \leq W \int_0^\infty \left(\int_0^y \left|\tilde{\Delta}_t^2 f^{(n)}\right|(x) \frac{(y-t)^{n-1}}{(n-1)!} dt\right) \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^\beta} dy. \tag{57}$$

Here we estimate in L_p norm, $p \geq 1$, the error function $K(x)$. Notice that we have $\omega_2(f^{(n)}, h)_p < \infty$, $h > 0$, $n = 0$ or $n \geq 2$ even. Operators M_ξ are positive operators.

The related main L_p result here comes next.

Theorem 3. *Let $p, q > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$, $n \geq 2$ even, $\alpha \in \mathbb{N}$, $\beta > \frac{1}{\alpha}\left(\frac{1}{p} + n + 2\right)$ and the rest as above. Then*

$$\|K(x)\|_p \leq \frac{\tilde{\tau}^{1/p} \alpha^{1/p} \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{1/q}}{2^{\frac{1}{q}} \Gamma\left(\frac{1}{2\alpha}\right)^{1/p} \Gamma\left(\beta - \frac{1}{2\alpha}\right) \Gamma\left(\frac{q\beta}{2}\right)^{1/q} (q(n-1)+1)^{1/q} (2p+1)^{1/p}} \frac{\Gamma(\beta)}{(n-1)!} \xi^n \omega_2(f^{(n)}, \xi)_p, \tag{58}$$

where

$$0 < \tilde{\tau} = \int_0^\infty \left((1+u)^{2p+1} - 1\right) u^{pn-1} \frac{1}{(1+u^{2\alpha})^{p\beta/2}} du < \infty. \tag{59}$$

Hence as $\xi \to 0$ we get $\|K(x)\|_p \to 0$.

If additionally $f^{(2m)} \in L_p(\mathbb{R}), m = 1, 2, \ldots, \frac{n}{2}$ then $\|M_\xi(f) - f\|_p \to 0$, as $\xi \to 0$.

Proof. We observe that

$$|K(x)|^p \leq W^p \left(\int_0^\infty \left(\int_0^y \left|\tilde{\Delta}_t^2 f^{(n)}\right|(x) \frac{(y-t)^{n-1}}{(n-1)!} dt\right) \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^\beta} dy\right)^p. \tag{60}$$

Call

$$\tilde{\gamma}(y,x) := \int_0^y |\tilde{\Delta}_t^2 f^{(n)}(x)| \frac{(y-t)^{n-1}}{(n-1)!} dt \geq 0, \quad y \geq 0, \tag{61}$$

then we have

$$|K(x)|^p \leq W^p \left(\int_0^\infty \tilde{\gamma}(y,x) \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^\beta} dy \right)^p . \tag{62}$$

Consequently

$$\begin{aligned}
\Lambda &:= \int_{-\infty}^{\infty} |K(x)|^p dx \leq W^p \int_{-\infty}^{\infty} \left(\int_0^\infty \tilde{\gamma}(y,x) \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^\beta} dy \right)^p dx \\
&= W^p \int_{-\infty}^{\infty} \left(\int_0^\infty \tilde{\gamma}(y,x) \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{\beta/2}} \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{\beta/2}} dy \right)^p dx \\
&\qquad \text{(by Hölder's inequality)} \\
&\leq W^p \left(\int_{-\infty}^{\infty} \left(\int_0^\infty (\tilde{\gamma}(y,x))^p \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{p\beta/2}} dy \right) \left(\int_0^\infty \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{q\beta/2}} dy \right)^{p/q} dx \right) \\
&= W^p \left(\frac{\Gamma\left(\frac{1}{2\alpha}\right) \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)}{2\Gamma\left(\frac{q\beta}{2}\right) \alpha \xi^{q\alpha\beta - 1}} \right)^{p/q} \left(\int_{-\infty}^{\infty} \left(\int_0^\infty (\tilde{\gamma}(y,x))^p \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{p\beta/2}} dy \right) dx \right) \\
&= \frac{\left[\Gamma\left(\beta\right)\right]^p \alpha \xi^{\alpha\beta p - 1} \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}} \Gamma\left(\frac{1}{2\alpha}\right) \Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{p/q}} \left(\int_{-\infty}^{\infty} \left(\int_0^\infty (\tilde{\gamma}(y,x))^p \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{p\beta/2}} dy \right) dx \right) \\
&=: (*).
\end{aligned} \tag{63}$$

By applying again Hölder's inequality we see that

$$\tilde{\gamma}(y,x) \leq \frac{\left(\int_0^y |\tilde{\Delta}_t^2 f^{(n)}(x)|^p dt \right)^{1/p}}{(n-1)!} \frac{y^{\left(n-1+\frac{1}{q}\right)}}{(q(n-1)+1)^{1/q}} . \tag{64}$$

Therefore it holds

$$\begin{aligned}
(*) &\leq \frac{\left[\Gamma\left(\beta\right)\right]^p \alpha \xi^{\alpha\beta p - 1} \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}} \Gamma\left(\frac{1}{2\alpha}\right) \Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{p/q} \left[(n-1)!\right]^p (q(n-1)+1)^{p/q}} \\
&\quad \cdot \left(\int_{-\infty}^{\infty} \left(\int_0^\infty \left(\int_0^y |\tilde{\Delta}_t^2 f^{(n)}(x)|^p dt \right) y^{p\left(n-1+\frac{1}{q}\right)} \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{p\beta/2}} dy \right) dx \right) \\
&= \frac{\left[\Gamma\left(\beta\right)\right]^p \alpha \xi^{\alpha\beta p - 1} \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}} \Gamma\left(\frac{1}{2\alpha}\right) \Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{p/q} \left[(n-1)!\right]^p (q(n-1)+1)^{p/q}} \\
&\quad \cdot \left(\int_0^\infty \left(\int_{-\infty}^{\infty} \left(\int_0^y |\tilde{\Delta}_t^2 f^{(n)}(x)|^p dt \right) y^{p\left(n-1+\frac{1}{q}\right)} \frac{1}{\left(y^{2\alpha} + \xi^{2\alpha}\right)^{p\beta/2}} dx \right) dy \right) \\
&=: (**).
\end{aligned} \tag{65}$$

We call

$$c_2 := \frac{\left[\Gamma\left(\beta\right)\right]^p \alpha \xi^{\alpha\beta p - 1} \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}} \Gamma\left(\frac{1}{2\alpha}\right) \Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{p/q} \left[(n-1)!\right]^p (q(n-1)+1)^{p/q}} . \tag{66}$$

And hence

$$
\begin{aligned}
(**) &= c_2 \left(\int_0^\infty \left(\int_{-\infty}^\infty \left(\int_0^y |\tilde{\Delta}_t^2 f^{(n)}(x)|^p dt \right) dx \right) y^{p(n-1+\frac{1}{q})} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right) \\
&= c_2 \left(\int_0^\infty \left(\int_0^y \left(\int_{-\infty}^\infty |\tilde{\Delta}_t^2 f^{(n)}(x)|^p dx \right) dt \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right) \\
&= c_2 \left(\int_0^\infty \left(\int_0^y \left(\int_{-\infty}^\infty |\Delta_t^2 f^{(n)}(x-t)|^p dx \right) dt \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right) \\
&= c_2 \left(\int_0^\infty \left(\int_0^y \left(\int_{-\infty}^\infty |\Delta_t^2 f^{(n)}(x)|^p dx \right) dt \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right) \\
&\le c_2 \left(\int_0^\infty \left(\int_0^y \omega_2(f^{(n)}, t)_p^p dt \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right) \\
&\le c_2 \omega_2(f^{(n)}, \xi)_p^p \left(\int_0^\infty \left(\int_0^y \left(1+\frac{t}{\xi}\right)^{2p} dt \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right). \qquad (67)
\end{aligned}
$$

I.e. so far we proved that

$$
\Lambda \le c_2 \omega_2(f^{(n)}, \xi)_p^p \left(\int_0^\infty \left(\int_0^y \left(1+\frac{t}{\xi}\right)^{2p} dt \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right). \qquad (68)
$$

But

$$
\text{R.H.S.}(68) = \frac{c_2 \xi}{2p+1} \omega_2(f^{(n)}, \xi)_p^p \left(\int_0^\infty \left(\left(1+\frac{y}{\xi}\right)^{2p+1} - 1 \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy \right). \qquad (69)
$$

Call

$$
M := \int_0^\infty \left(\left(1+\frac{y}{\xi}\right)^{2p+1} - 1 \right) y^{pn-1} \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{p\beta/2}} dy, \qquad (70)
$$

and

$$
\tilde{\tau} := \int_0^\infty \left((1+u)^{2p+1} - 1 \right) u^{pn-1} \frac{1}{(1+u^{2\alpha})^{p\beta/2}} du. \qquad (71)
$$

That is

$$
M = \xi^{p(n-\alpha\beta)} \tilde{\tau}. \qquad (72)
$$

Therefore it holds

$$
\Lambda \le \frac{\tilde{\tau} \left[\Gamma(\beta)\right]^p \alpha \xi^{pn} \Gamma\left(\frac{q\beta}{2} - \frac{1}{2\alpha}\right)^{p/q} \omega_2(f^{(n)}, \xi)_p^p}{2^{\frac{p}{q}} (2p+1) \Gamma\left(\frac{1}{2\alpha}\right) \Gamma\left(\beta - \frac{1}{2\alpha}\right)^p \Gamma\left(\frac{q\beta}{2}\right)^{p/q} [(n-1)!]^p (q(n-1)+1)^{p/q}}. \qquad (73)
$$

We have established (58). ■

The counterpart of Theorem 3 follows, $p = 1$ case.

Theorem 4. *Let* $f \in C^n(\mathbb{R})$ and $f^{(n)} \in L_1(\mathbb{R})$, $n \geq 2$ *even*, $\alpha \in \mathbb{N}$, $\beta > \frac{n+3}{2\alpha}$. *Then*

$$\|K(x)\|_1 \leq \frac{1}{6\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta - \frac{1}{2\alpha}\right)(n-1)!}\left[3\Gamma\left(\frac{n+1}{2\alpha}\right)\Gamma\left(\beta - \frac{n+1}{2\alpha}\right)\right. \tag{74}$$

$$\left.+3\Gamma\left(\frac{n+2}{2\alpha}\right)\Gamma\left(\beta - \frac{n+2}{2\alpha}\right) + \Gamma\left(\frac{n+3}{2\alpha}\right)\Gamma\left(\beta - \frac{n+3}{2\alpha}\right)\right]\omega_2(f^{(n)},\xi)_1\xi^n.$$

Hence as $\xi \to 0$ *we obtain* $\|K(x)\|_1 \to 0$.

If additionally $f^{(2m)} \in L_1(\mathbb{R}), m = 1, 2, \ldots, \frac{n}{2}$ then $\|M_\xi(f) - f\|_1 \to 0$, as $\xi \to 0$.

Proof. Notice that

$$\tilde{\Delta}_t^2 f^{(n)}(x) = \Delta_t^2 f^{(n)}(x-t), \tag{75}$$

all $x, t \in \mathbb{R}$. Also it holds

$$\int_{-\infty}^{\infty} |\Delta_t^2 f^{(n)}(x-t)|dx = \int_{-\infty}^{\infty} |\Delta_t^2 f^{(n)}(w)|dw \leq \omega_2(f^{(n)}, t)_1, \quad \text{all } t \in \mathbb{R}_+. \tag{76}$$

Here we obtain

$$\begin{aligned}
\|K(x)\|_1 &= \int_{-\infty}^{\infty} |K(x)|dx \\
&\overset{(57)}{\leq} W\int_{-\infty}^{\infty}\left(\int_0^{\infty}\left(\int_0^y \left|\tilde{\Delta}_t^2 f^{(n)}(x)\right| \frac{(y-t)^{n-1}}{(n-1)!}dt\right)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dy\right)dx \\
&\leq W\int_{-\infty}^{\infty}\left(\int_0^{\infty}\left(\frac{y^{n-1}}{(n-1)!}\int_0^y \left|\tilde{\Delta}_t^2 f^{(n)}(x)\right| dt\right)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dy\right)dx \\
&= W\int_0^{\infty}\left(\left(\int_{-\infty}^{\infty}\left(\int_0^y \left|\tilde{\Delta}_t^2 f^{(n)}(x)\right| dt\right)dx\right)\frac{y^{n-1}}{(n-1)!}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}\right)dy \\
&\overset{(75)}{=} W\int_0^{\infty}\left(\left(\int_{-\infty}^{\infty}\left(\int_0^y \left|\Delta_t^2 f^{(n)}(x-t)\right| dt\right)dx\right)\frac{y^{n-1}}{(n-1)!}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}\right)dy \\
&= W\int_0^{\infty}\left(\left(\int_0^y\left(\int_{-\infty}^{\infty} \left|\Delta_t^2 f^{(n)}(x-t)\right| dx\right)dt\right)\frac{y^{n-1}}{(n-1)!}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}\right)dy \\
&\overset{(76)}{\leq} W\int_0^{\infty}\left(\left(\int_0^y \omega_2(f^{(n)}, t)_1 dt\right)\frac{y^{n-1}}{(n-1)!}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}\right)dy \\
&\leq W\omega_2(f^{(n)},\xi)_1\left(\int_0^{\infty}\left(\int_0^y\left(1+\frac{t}{\xi}\right)^2 dt\right)\frac{y^{n-1}}{(n-1)!}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dy\right) \\
&= W\omega_2(f^{(n)},\xi)_1\left(\int_0^{\infty}\left(\left(1+\frac{y}{\xi}\right)^3 - 1\right)\frac{\xi}{3}\frac{y^{n-1}}{(n-1)!}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta}}dy\right) \\
&= \frac{\Gamma(\beta)\,\alpha\xi^n}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)(n-1)!3}\omega_2(f^{(n)},\xi)_1\left(\int_0^{\infty}\left((1+Y)^3-1\right)Y^{n-1}\frac{1}{\left(Y^{2\alpha}+1\right)^{\beta}}dY\right)
\end{aligned}$$

$$= \frac{\Gamma(\beta)\alpha\xi^n}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)(n-1)!3}\omega_2(f^{(n)},\xi)_1\left(\int_0^\infty \left(3Y+3Y^2+Y^3\right)Y^{n-1}\frac{1}{\left(Y^{2\alpha}+1\right)^\beta}dY\right)$$

$$= \frac{\Gamma(\beta)\alpha\xi^n}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)(n-1)!3}\omega_2(f^{(n)},\xi)_1\left(\int_0^\infty \left(3Y^n+3Y^{n+1}+Y^{n+2}\right)\frac{1}{\left(Y^{2\alpha}+1\right)^\beta}dY\right)$$

$$= \frac{1}{6\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)(n-1)!}\left[3\Gamma\left(\frac{n+1}{2\alpha}\right)\Gamma\left(\beta-\frac{n+1}{2\alpha}\right)\right.$$
$$\left.+3\Gamma\left(\frac{n+2}{2\alpha}\right)\Gamma\left(\beta-\frac{n+2}{2\alpha}\right)+\Gamma\left(\frac{n+3}{2\alpha}\right)\Gamma\left(\beta-\frac{n+3}{2\alpha}\right)\right]\omega_2(f^{(n)},\xi)_1\xi^n. \tag{77}$$

We have proved (74). ■

The related case here of $n=0$ comes next.

Proposition 3. *Let $p,q>1$ such that $\frac{1}{p}+\frac{1}{q}=1$, $\alpha\in\mathbb{N}$, $\beta>\frac{1}{\alpha}\left(2+\frac{1}{p}\right)$ and the rest as above. Then*

$$\|M_\xi(f)-f\|_p \le \frac{\rho^{1/p}\Gamma(\beta)\alpha^{1/p}\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{1/q}}{2^{\frac{1}{q}}\Gamma\left(\frac{1}{2\alpha}\right)^{1/p}\Gamma\left(\beta-\frac{1}{2\alpha}\right)\Gamma\left(\frac{q\beta}{2}\right)^{1/q}}\omega_2(f,\xi)_p, \tag{78}$$

where

$$0<\rho:=\int_0^\infty (1+y)^{2p}\frac{1}{\left(y^{2\alpha}+1\right)^{\beta p/2}}dy<\infty. \tag{79}$$

Hence as $\xi\to 0$ we obtain $M_\xi\to I$ in the L_p norm, $p>1$.

Proof. From (55) we get

$$|M_\xi(f;x)-f(x)|^p \le W^p\left(\int_0^\infty \left|\tilde{\Delta}_y^2 f(x)\right|\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy\right)^p. \tag{80}$$

We then estimate

$$\int_{-\infty}^\infty |M_\xi(f;x)-f(x)|^p dx \le W^p\int_{-\infty}^\infty\left(\int_0^\infty \left|\tilde{\Delta}_y^2 f(x)\right|\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy\right)^p dx$$

$$= W^p\int_{-\infty}^\infty\left(\int_0^\infty \left|\tilde{\Delta}_y^2 f(x)\right|\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}dy\right)^p dx$$

$$\le W^p\int_{-\infty}^\infty\left(\left(\int_0^\infty\left(\left|\tilde{\Delta}_y^2 f(x)\right|\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right)^p dy\right)^{\frac{1}{p}}\left(\int_0^\infty\left(\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta/2}}\right)^q dy\right)^{\frac{1}{q}}\right)^p dx$$

$$= W^p\int_{-\infty}^\infty\left(\left(\int_0^\infty\left|\tilde{\Delta}_y^2 f(x)\right|^p\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta p/2}}dy\right)\left(\int_0^\infty\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta q/2}}dy\right)^{\frac{p}{q}}\right)dx$$

$$= W^p\left(\int_0^\infty\left(\int_{-\infty}^\infty\left|\tilde{\Delta}_y^2 f(x)\right|^p\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta p/2}}dx\right)dy\right)\left(\frac{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)}{2\Gamma\left(\frac{q\beta}{2}\right)\alpha\xi^{q\alpha\beta-1}}\right)^{\frac{p}{q}}$$

$$
\begin{aligned}
&= \frac{[\Gamma(\beta)]^p \alpha \xi^{\alpha\beta p-1}\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p\Gamma\left(\frac{q\beta}{2}\right)^{p/q}} \\
&\cdot\left(\int_0^\infty\left(\int_{-\infty}^\infty \left|\Delta_y^2 f(x-y)\right|^p dx\right)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta p/2}}dy\right) \\
&\le \frac{[\Gamma(\beta)]^p \alpha \xi^{\alpha\beta p-1}\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p\Gamma\left(\frac{q\beta}{2}\right)^{p/q}}\int_0^\infty \omega_2(f,y)_p^p\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta p/2}}dy \\
&\le \frac{[\Gamma(\beta)]^p \alpha \xi^{\alpha\beta p-1}\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p\Gamma\left(\frac{q\beta}{2}\right)^{p/q}}\omega_2(f,\xi)_p^p\int_0^\infty\left(1+\frac{y}{\xi}\right)^{2p}\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^{\beta p/2}}dy \\
&= \frac{[\Gamma(\beta)]^p \alpha \xi^{\alpha\beta p-1}\Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p\Gamma\left(\frac{q\beta}{2}\right)^{p/q}}\omega_2(f,\xi)_p^p\int_0^\infty(1+Y)^{2p}\frac{1}{\left(Y^{2\alpha}+1\right)^{\beta p/2}\xi^{\alpha\beta p}}\xi dY \\
&= \frac{[\Gamma(\beta)]^p \alpha \Gamma\left(\frac{q\beta}{2}-\frac{1}{2\alpha}\right)^{p/q}}{2^{\frac{p}{q}}\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)^p\Gamma\left(\frac{q\beta}{2}\right)^{p/q}}\omega_2(f,\xi)_p^p\int_0^\infty(1+y)^{2p}\frac{1}{\left(y^{2\alpha}+1\right)^{\beta p/2}}dy. \qquad (81)
\end{aligned}
$$

The proof of (78) is now completed. ■

Also we give

Proposition 4. *For $\alpha\in\mathbb{N}$, $\beta>\frac{3}{2\alpha}$, it holds,*

$$
\|M_\xi f-f\|_1\le\left[\frac{1}{2}+\frac{\Gamma\left(\frac{1}{\alpha}\right)\Gamma\left(\beta-\frac{1}{\alpha}\right)}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}+\frac{\Gamma\left(\frac{3}{2\alpha}\right)\Gamma\left(\beta-\frac{3}{2\alpha}\right)}{2\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}\right]\omega_2(f,\xi)_1. \qquad (82)
$$

Hence as $\xi\to 0$ we get $M_\xi\to I$ in the L_1 norm.

Proof. From (55) we have

$$
|M_\xi(f;x)-f(x)|\le W\left(\int_0^\infty\left|\tilde{\Delta}_y^2 f(x)\right|\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy\right). \qquad (83)
$$

Hence we get

$$
\begin{aligned}
\int_{-\infty}^\infty|M_\xi(f;x)-f(x)|dx &\le W\int_{-\infty}^\infty\left(\int_0^\infty|\tilde{\Delta}_y^2 f(x)|\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy\right)dx \\
&= W\int_0^\infty\left(\int_{-\infty}^\infty|\tilde{\Delta}_y^2 f(x)|dx\right)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy \\
&= W\int_0^\infty\left(\int_{-\infty}^\infty|\Delta_y^2 f(x-y)|dx\right)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy \\
&= W\int_0^\infty\left(\int_{-\infty}^\infty|\Delta_y^2 f(x)|dx\right)\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy \\
&\le W\int_0^\infty\omega_2(f,y)_1\frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta}dy
\end{aligned}
$$

$$
\begin{aligned}
&\leq W\omega_2(f,\xi)_1 \int_0^\infty \left(1+\frac{y}{\xi}\right)^2 \frac{1}{\left(y^{2\alpha}+\xi^{2\alpha}\right)^\beta} dy \\
&= W\omega_2(f,\xi)_1 \int_0^\infty (1+x)^2 \frac{1}{\left(x^{2\alpha}+1\right)^\beta \xi^{2\alpha\beta}} \xi dx \\
&= \frac{\Gamma(\beta)\alpha}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)} \omega_2(f,\xi)_1 \int_0^\infty (1+x)^2 \frac{1}{\left(x^{2\alpha}+1\right)^\beta} dx \\
&= \left[\frac{1}{2} + \frac{\Gamma\left(\frac{1}{\alpha}\right)\Gamma\left(\beta-\frac{1}{\alpha}\right)}{\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)} + \frac{\Gamma\left(\frac{3}{2\alpha}\right)\Gamma\left(\beta-\frac{3}{2\alpha}\right)}{2\Gamma\left(\frac{1}{2\alpha}\right)\Gamma\left(\beta-\frac{1}{2\alpha}\right)}\right] \omega_2(f,\xi)_1. \qquad (84)
\end{aligned}
$$

We have established (82). ■

References

[1] G.A. Anastassiou, Rate of convergence of non-positive linear convolution type operators. A sharp inequality, *J. Math. Anal. and Appl.*, **142** (1989), 441–451.

[2] G.A. Anastassiou, Sharp inequalities for convolution type operators, *Journal of Approximation Theory*, **58** (1989), 259–266.

[3] G.A. Anastassiou, *Moments in Probability and Approximation Theory*, Pitman Research Notes in Math., Vol. 287, Longman Sci. & Tech., Harlow, U.K., 1993.

[4] G.A. Anastassiou, *Quantitative Approximations*, Chapman & Hall/CRC, Boca Raton, New York, 2001.

[5] G.A. Anastassiou, Basic convergence with rates of smooth Picard singular integral operators, *J. of Computational Analysis and Applications*,Vol.8, No.4 (2006), 313-334.

[6] G.A. Anastassiou, L_p convergence with rates of smooth Picard singular operators, *Differential & difference equations and applications, Hindawi Publ. Corp., New York,* (2006), 31–45.

[7] G.A. Anastassiou and S. Gal, Convergence of generalized singular integrals to the unit, univariate case, *Math. Inequalities & Applications*, **3**, No. 4 (2000), 511–518.

[8] G.A. Anastassiou and S. Gal, Convergence of generalized singular integrals to the unit, multivariate case, *Applied Math. Rev.*, Vol. 1, World Sci. Publ. Co., Singapore, 2000, pp. 1–8.

[9] G.A. Anastassiou and R. Mezei, *L_p Convergence with Rates of Smooth Gauss-Weierstrass Singular Operators*, Nonlinear Studies, accepted 2008.

[10] G.A. Anastassiou and R. A. Mezei, *Global Smoothness and Uniform Convergence of Smooth Poisson-Cauchy Type Singular Operators,* submitted 2009.

[11] G.A. Anastassiou and R. A. Mezei, *A Voronovskaya Type Theorem for Poisson-Cauchy Type Singular Operators,* submitted 2009.

[12] R.A. DeVore and G.G. Lorentz, *Constructive Approximation*, Springer-Verlag, Vol. 303, Berlin, New York, 1993.

[13] S.G. Gal, Remark on the degree of approximation of continuous functions by singular integrals, *Math. Nachr.*, **164** (1993), 197–199.

[14] S.G. Gal, Degree of approximation of continuous functions by some singular integrals, *Rev. Anal. Numér, Théor. Approx.*, (Cluj), Tome XXVII, No. 2 (1998), 251–261.

[15] R.N. Mohapatra and R.S. Rodriguez, On the rate of convergence of singular integrals for Hölder continuous functions, *Math. Nachr.* **149** (1990), 117–124.

[16] D. Zwillinger, *CRC Standard Mathematical Tables and Formulae, 30th Edition,* Chapman & Hall/CRC, Boca Raton, 1995.